


Product Glossary

Affordability Index of Biocon Products



Cardiology



STATIX®

Active Ingredient: Atorvastatin
10/20/40 mg
Indication: Controls elevated cholesterol levels



STATIX®-F

Active Ingredient: Atorvastatin
10 mg + Micronised Fenofibrate
200 mg
Indication: Corrects cholesterol levels in Diabetic Dyslipidemics



STATIX® - EZ

Active Ingredient: Atorvastatin
10 mg + Ezetimibe 10 mg
Indication: Controls extremely high levels of cholesterol



TELMISAT®

Active Ingredient: Telmisartan
20/40/80 mg tablets
Indication: Offers 24 hour blood pressure control



TELMISAT® -H

Active Ingredient: Telmisartan
40/80 mg + Hydrochlorothiazide
12.5 mg
Indication: In uncontrolled hypertension



TELMISAT® AM

Active Ingredient: Telmisartan
40 mg + Amlodipine 5 mg tablets
Indication: In Diabetic Hypertensives



ACTIBLOK™ - IPR

Active Ingredient: Metoprolol
Immediate & Patterned Release
25/50/100 mg tablets
Indication: In patients of Hypertension, Angina, IHD and Heart Failure



ACTIBLOK™ AM

Active Ingredient: Metoprolol
Succinate IPR 25/50 mg +
Amlodipine 5 mg tablets
Indication: In Uncontrolled Hypertension



BESTOR®

Active Ingredient: Rosuvastatin
Calcium 5/10 mg tablets
Indication: For the management of Dyslipidemia and Atherosclerosis



BRADIA™

Active Ingredient: Ivabradine
5/7.5 mg tablets
Indication: For the management of Stable Angina



THINRIN™
Active Ingredient: Clopidogrel
 75/150 mg
Indication: For early and long term risk reduction in high risk ACS patients



CLASPRIN®
Active Ingredient: Aspirin
 75/150 mg + Clopidogrel 75 mg
Indication: For early and long term risk reduction in high risk ACS patients



ZARGO®
Active Ingredient: Losartan Potassium 25/50 mg
Indication: Reduces high blood pressure



ZARGO® - H
Active Ingredient: Losartan Potassium 50 mg + Hydrochlorothiazide 12.5 mg
Indication: Reduces high blood pressure



ZIGPRIL®
Active Ingredient: Ramipril 2.5/5 mg
Indication: In patients with risk of CVD



MYOKINASE™
Active Ingredient: Recombinant Streptokinase for injection 1,500,000 IU
Indication: In patients of Acute Myocardial Infarction



DYNALIX®
Active Ingredient: Enoxaparin 40 mg, 60 mg Pre Filled Syringe
Indication: In patients of Acute Coronary Syndrome and Prophylaxis of Deep Vein Thrombosis



CLOTIDE™
Active Ingredient: Eptifi batide 10 ml bolus, 100 ml infusion
Indication: In patients of Acute Coronary Syndrome, undergoing Percutaneous Coronary Interventions

Diabetology



INSUGEN® -R (Regular)
INSUGEN® -N(NPH)
INSUGEN® -30/70 & 50/50 (Biphasic)
Active Ingredient: Each ml contains Human Insulin (rDNA origin), Ph Eur 40 IU
Indication: In Diabetes, useful when oral agents fail to control blood glucose levels



BASALOG™
Active Ingredient: Each ml contains Insulin Glargine (rDNA Origin) 100 IU
Indication: In Diabetes Mellitus, for 24 hrs basal insulin action



BLISTO™
Active Ingredient: Glimepiride
 1/2/4 mg
Indication: Increases Insulin secretion in Type 2 Diabetes by stimulating pancreas



BLISTO™ -1MF/2MF/4MF
Active Ingredient: Glimepiride 1 mg + Metformin 500 mg SR & Glimepiride 2/4 mg + Metformin 1000 mg SR
Indication: In Type 2 Diabetes, when blood glucose is not controlled with a single medication



METADOZE®-IPR
Active Ingredient: Metformin
 500/850 mg SR
Indication: Improves action of Insulin in Type 2 Diabetes



TriGPM™-1/2
Active Ingredient: Glimepiride 1/2 mg + Pioglitazone 15 mg + Metformin 500 mg SR
Indication: A triple drug combination for uncontrolled Type 2 Diabetes



ZUKER™-MF
Active Ingredient: Gliclazide 80 mg + Metformin 500 mg SR
Indication: Comprehensively controls hyperglycemia in Type 2 Diabetes



PIODART®
Active Ingredient: Pioglitazone 15/30 mg
Indication: Helps Insulin work better in Type 2 Diabetes



PIODART®-MF
Active Ingredient: Pioglitazone 15 mg + Metformin ER 500 mg
Indication: Improves blood sugar control when not controlled by monotherapy



OLISAT™
Active Ingredient: Orlistat 60/120 mg
Indication: Weight reducer



GABIL™
Active Ingredient: Gabapentin 300 mg + Methylcobalamin 500 mcg
Indication: In Diabetic Neuropathy



GMAB™ Plus
Active Ingredient: GLA 100 mg + Methylcobalamin 1500 mcg + ALA 100 mg + Benfothiamine 100 mg + Ele. Zinc 15 mg
Indication: Nutritional supplement

Nephrology



ERYPRO safe™

Active Ingredient: Recombinant Human Erythropoietin Alpha injection in strengths of 2000 IU/3000 IU/4000 IU/5000 IU/6000 IU/10000 IU
Indication: For the treatment of patients with anemia due to chronic renal failure, either on dialysis or non-dialysis



ERYPRO™

Active Ingredient: Recombinant Human Erythropoietin Alpha 2000 IU/4000 IU/10000 IU
Indication: For the treatment of patients with anemia due to chronic renal failure, either on dialysis or non-dialysis



TACROGRAF™

Active Ingredient: Tacrolimus 0.5/1/2/3/5 mg capsules
Indication: Prophylaxis of transplant rejection in organ transplantation and as a rescue therapy in patients with rejection episodes



RENODAPT®-S

Active Ingredient: Mycophenolic Acid 180/360/540 mg tablets
Indication: Prophylaxis of transplant rejection in organ transplantation and as a rescue therapy in patients with rejection episodes



RENODAPT®

Active Ingredient: Mycophenolate Mofetil 250 mg capsules and 500/750 mg tablets
Indication: Prophylaxis of transplant rejection in organ transplantation and as a rescue therapy in patients with rejection episodes



CYCLOPHIL ME™

Active Ingredient: Cyclosporine USP 25/50/100 mg capsules
Indication: Prophylaxis of allograft rejection in kidney transplantation and as a rescue therapy in patients with rejection episodes



CYCLOPHIL ME™ (ORAL SOLUTION)

Active Ingredient: Cyclosporine Oral Solution USP 100 mg/ml
Indication: Prophylaxis of transplant rejection in organ transplantation and as a rescue therapy in patients with rejection episodes



Narita™+

Active Ingredient: Whey protein supplement fortified with vitamins and minerals, 200 gm tin
Indication: For patients who need high protein supplementation



RAPACAN™

Active Ingredient: Sirolimus 1/2 mg tablets
Indication: Prevention of rejection and rescue therapy for rejection in renal transplantation



bioSEV™

Active Ingredient: Sevelamer HCl 400/800 mg tablets
Indication: For the management of hyperphosphatemia in CKD patients



CeRACal™

Active Ingredient: Cinacalcet hydrochloride equivalent to Cinacalcet 30/60 mg

Indication: For the treatment of secondary hyperparathyroidism in dialysis patients

Oncology



BIOMAb EGFR®

Active Ingredient: Nimotuzumab 200 mg

Indication: Humanized monoclonal antibody targeting epidermal growth factor receptor indicated for its use in head and neck cancer



Abraxane®

Active Ingredient: Paclitaxel protein bound particles for injectable suspension (Albumin bound)

Indication: For the treatment of Breast Cancer after failure of combination chemotherapy for metastatic disease or relapse within 6 months of adjuvant chemotherapy



ERYPRO safe™

Active Ingredient: Recombinant Human Erythropoietin Alpha 10000 IU/40000 IU

Indication: For the treatment of chemotherapy induced anemia



NUFIL safe™

Active Ingredient: Filgrastim (Recombinant Human Granulocyte Colony Stimulating Factor) 300 µg

Indication: For the treatment of chemotherapy induced neutropenia



NUFIL™

Active Ingredient: Filgrastim (Recombinant Human Granulocyte Colony Stimulating Factor) 300 µg

Indication: For the treatment of chemotherapy induced neutropenia