


PRODUCT GLOSSARY

Affordability Index of Biocon Products



Cardiology



STATIX®

Active Ingredient: Atorvastatin
10/20/40/80 mg tablets
Indication: Controls elevated cholesterol levels



STATIX® F

Active Ingredient: Atorvastatin
10 mg + Micronised Fenofibrate
200 mg tablets
Indication: For Diabetic Dyslipidemia



STATIX® - EZ

Active Ingredient: Atorvastatin
10 mg + Ezetimibe 10 mg tablets
Indication: Controls extremely high levels of cholesterol



TELMISAT®

Active Ingredient: Telmisartan
20/40/80 mg tablets
Indication: Offers 24 hour blood pressure control



TELMISAT®-H

Active Ingredient: Telmisartan
40/80 mg + Hydrochlorothiazide
12.5 mg tablets
Indication: In uncontrolled Hypertension



TELMISAT® AM

Active Ingredient: Telmisartan
40 mg + Amlodipine 5 mg tablets
Indication: In Diabetic Hypertensives



ACTIBLOK™ - IPR

Active Ingredient: Metoprolol
Immediate & Patterned Release
25/50/100 mg tablets
Indication: In patients of Hypertension, Angina, IHD and Heart Failure



ACTIBLOK AM™

Active Ingredient: Metoprolol
Succinate IPR 25/50 mg +
Amlodipine 5 mg tablets
Indication: In Uncontrolled Hypertension




BESTOR®

Active Ingredient: Rosuvastatin
Calcium 5/10/20 mg tablets
Indication: For the management of Dyslipidemia and Atherosclerosis



BRADIA™

Active Ingredient: Ivabradine
5 mg tablets
Indication: For the management of Stable Angina



CLASPRIN®

Active Ingredient: Aspirin
75/150 mg + Clopidogrel 75 mg
capsules

Indication: For early and long
term risk reduction in high risk
ACS patients



ZARGO®

Active Ingredient: Losartan
Potassium 25/50 mg tablets

Indication: In hypertensive
patients with CV co-morbidity



ZARGO® - H

Active Ingredient: Losartan
Potassium 50 mg +
Hydrochlorothiazide 12.5 mg
tablets

Indication: In severe Hypertensive
patients



MYOKINASE®

Active Ingredient: Recombinant
Streptokinase for injection
1,500,000 IU

Indication: In patients of Acute
Myocardial Infarction



DYNALIX®

Active Ingredient: Enoxaparin
40/60 mg Pre Filled Syringe

Indication: In patients of Acute
Coronary Syndrome and Prophylaxis
of Deep Vein Thrombosis



CLOTIDE®

Active Ingredient: Eptifibatide
10 ml bolus/100 ml for infusion

Indication: In patients of Acute
Coronary Syndrome, undergoing
Percutaneous Coronary Interventions



PRASACT™

Active Ingredient: Prasugrel 5/10 mg
tablets

Indication: Reduces thrombotic CV
events



TIROZEST™

Active Ingredient: Tirozest HCL
5 mg/100 ml vial

Indication: In high risk ACS patients
with Acute MI

Diabetology



INSUGEN® 30/70 40 IU
INSUGEN® 50/50 40 IU
INSUGEN® N 40 IU
INSUGEN® R 40 IU

Active Ingredient: Each ml contains Human Insulin (rDNA origin), IP 40 IU
Indication: In Diabetes, useful when oral agents fail to control blood glucose levels



Insugen® 30/70 100 IU
Insugen® 50/50 100 IU
Insugen® N 100 IU
Insugen® R 100 IU

Active Ingredient: Each ml contains Human Insulin (rDNA origin), IP 100 IU
Indication: In Diabetes, useful when oral agents fail to control blood glucose levels




BASALOG®

Active Ingredient: Each ml contains Insulin Glargine (rDNA Origin) 100 IU
Indication: In Diabetes Mellitus, for 24 hrs basal insulin action



BLISTO®

Active Ingredient: Glimperide 1/2/4 mg
Indication: Oral antidiabetic agent that acts by stimulating beta cells leading to insulin secretion




BLISTO MF®

Active Ingredient: Glimperide 1 mg + Metformin 500 mg SR & Glimperide 2/4 mg + Metformin 1000 mg SR
Indication: Oral anti-diabetic agent that combination therapy that controls hyperglycaemia in Type 2 Diabetes




METADOZE-IPR®

Active Ingredient: Metformin 500/850 mg IPR
Indication: Oral antidiabetic agent that improves action of Insulin in Type 2 Diabetes



TriGPM®

Active Ingredient: Glimperide 1/2 mg + Pioglitazone 15 mg + Metformin 500 mg ER
Indication: A triple drug combination for the management of Type 2 diabetes, uncontrolled with combination therapy of two drugs



ZUKER®-MF

Active Ingredient: Gliclazide 80 mg + Metformin 500 mg SR
Indication: Oral anti-diabetic combination therapy for controlling hyperglycaemia in Type 2 diabetes




PIODART®

Active Ingredient: Pioglitazone 15/30 mg
Indication: Oral anti-diabetic agent that improves action of insulin in Type 2 Diabetes



PIODART®-MF

Active Ingredient: Pioglitazone 15 mg + Metformin ER 500 mg
Indication: Improves blood sugar control when hyperglycaemia is not controlled by monotherapy


**OLISAT®****Active Ingredient:**

Orlistat 60/120 mg

Indication: Helps in Weight reduction**GABIL®****Active Ingredient:** Gabapentin

300 mg + Methylcobalamin

500 mcg

Indication: For reduction in diabetic neuropathy symptoms**GMAB® Plus****Active Ingredient:** GLA 100 mg

+ Methylcobalamin 1500 mcg + ALA

100 mg + Benfothiamine 100 mg +

Elemental Zinc 15 mg

Indication: Nutritional supplement used along with other anti-diabetic medications

Nephrology

**ERYPRO safe™****Active Ingredient:** Recombinant Human Erythropoietin Alpha injection in strengths of 2000 IU/3000 IU/4000 IU/5000 IU/6000 IU/10000 IU**Indication:** For the treatment of patients with anemia due to chronic renal failure, either on dialysis or not on dialysis**ERYPRO™****Active Ingredient:** Recombinant Human Erythropoietin Alpha 2000 IU/4000 IU/10000 IU**Indication:** For the treatment of patients with anemia due to chronic renal failure, either on dialysis or not on dialysis**TACROGRAF™****Active Ingredient:** Tacrolimus 0.5/1/2/3/5 mg capsules**Indication:** Prophylaxis of organ rejection in patients receiving allogenic liver, kidney or heart transplantation**RENODAPT®****Active Ingredient:** Mycophenolate Mofetil 250 mg capsules and 500/750 mg tablets**Indication:** Prophylaxis of organ rejection in patients receiving allogenic renal, heart or liver transplants**RENODAPT®-S****Active Ingredient:** Mycophenolic Acid 180/360/540 mg tablets**Indication:** Prophylaxis of organ rejection in patients receiving allogenic renal transplants**CYCLOPHIL ME® (ORAL SOLUTION)****Active Ingredient:** Cyclosporine Oral Solution USP 100 mg/ml**Indication:** Prophylaxis of organ rejection in kidney, liver and heart allogenic transplants

**CYCLOPHIL ME®**

Active Ingredient: Cyclosporine
USP 25/50/100 mg capsules
Indication: Prophylaxis of organ rejection in kidney, liver and heart allogenic transplants

**RAPACAN™**

Active Ingredient: Sirolimus
1/2 mg tablets
Indication: Prophylaxis of organ rejection in patients aged 13yrs or older receiving renal transplants

**Narita™+**

Active Ingredient: Whey protein supplement fortified with vitamins and minerals, 200 gm tin
Indication: Used as a nutritional supplement in chronic kidney patients undergoing dialysis

**CeRACal™**

Active Ingredient: Cinacalcet hydrochloride equivalent to Cinacalcet 30/60 mg tablets
Indication: Treatment of secondary hyperparathyroidism in dialysis patients

**biOSeV™**

Active Ingredient: Sevelamer HCl 400/800 mg tablets
Indication: Control of serum phosphorous in patients with chronic kidney disease on dialysis

**Advacan™**

Active Ingredient: Everolimus 0.25/0.5 mg tablets
Indication: Prophylaxis of organ rejection in adult patients at low to moderate immunologic risk receiving a kidney or cardiac transplant

Oncotherapeutics

**BIOMab EGFR®**

Active Ingredient: Nimotuzumab 200 mg humanized monoclonal antibody targeting epidermal growth factor receptor
Indication: For the treatment of locally advanced squamous cell carcinoma of head and neck, along with radiation and/or chemotherapy

**Abraxane®**

Active Ingredient: Paclitaxel protein-bound particles for injectable suspension (albumin-bound)
Indication: Abraxane for Injectable Suspension is indicated for the treatment of breast cancer after failure of combination chemotherapy for metastatic disease or relapse within 6 months of adjuvant chemotherapy



ERYPRO safe™

Active Ingredient:

Recombinant Human Erythropoietin Alpha 10000 IU/40000 IU

Indication: For the treatment of chemotherapy induced anemia



NUFIL safe™

Active Ingredient: Filgrastim

(Recombinant Human Granulocyte Colony Stimulating Factor) 300 µg

Indication: For the treatment of chemotherapy induced neutropenia



NUFIL™

Active Ingredient: Filgrastim

(Recombinant Human Granulocyte Colony Stimulating Factor) 300 µg

Indication: For the treatment of chemotherapy induced neutropenia



Evertor™

Active Ingredient: Everolimus 5 mg/10 mg

Indication: For the treatment of patients with advanced renal cell carcinoma (RCC) after failure of treatment with Sunitinib or Sorafenib

Immunology



PICON®

Active Ingredient: Pimecrolimus cream 1% w/w

Indication: Mild to Moderate atopic Dermatitis



TBIS®

Active Ingredient: Tacrolimus ointment 0.03%/0.1% w/w

Indication: Moderate to Severe Dermatitis

Comprehensive Care



CELRIM®

Active Ingredient: Cefepime 1gr

Indication: • Pneumonia • Febrile Neutropenia • Urinary Tract Infections • Uncomplicated Skin and Skin Structure Infections • Complicated Intra-abdominal Infections



MEEZAT™

Active Ingredient: Ceftazidime 1g

Indication: • Lower Respiratory Tract Infections • Skin and Skin-Structure Infections • Urinary Tract Infections • Bacterial Septicemia • Bone and Joint Infections • Gynecologic Infections • Intra-abdominal Infections • Central Nervous System Infections

**Biopiper TZ™****Active Ingredient:**

Piperacillin 4 g & Tazobactam 0.5 g

Indication: • Intra- abdominal Infections • Skin and skin structure infections • Postpartum endometritis or pelvic inflammatory disease • Community-acquired pneumonia**CELRIM TZ®****Active Ingredient:** Cefepime 1g & Tazo bactam 0.125 g**Indication:** • Urinary tract infections • Skin and skin structure infections and • Complicated intra-abdominal infections**IMICELUM™****Active Ingredient:** Imipenem 500 mg & Cilastatin 500 mg**Indication:** • Lower respiratory tract infections • Urinary tract infections • Intra-abdominal infections • Gyneacologic infections • Bacterial septicemia • Bone and joint infections • Skin and skin structure infections • Endocarditis**ENTAVAR™ – 600****Active Ingredient:** Linezolid Tablet 600 mg**Indication:** In Gram-Positive infections • VRE infections • Nosocomial pneumonia • Skin and skin structure infections • Diabetic foot infections**PENMER® – 500****Active Ingredient:** Meropenem 500 mg**Indication:** • Skin and Skin Structure Infections • Intra-abdominal Infections • Bacterial Meningitis**GENPIROME®****Active Ingredient:** Cefpirome 1g**Indication:** • Lower respiratory tract infections • Complicated upper and lower urinary tract infections • Bacteremia/ Septicaemia • Febrile neutropenia • Skin and soft tissue infections**ENTAVAR™****Active Ingredient:** Linezolid I.V 200 mg/300 ml**Indication:** In Gram-Positive infections: • VRE infections • Nosocomial pneumonia • Skin and skin structure infections • Diabetic foot infections**PENMER®****Active Ingredient:** Meropenem 1g**Indication:** • Skin and Skin Structure Infections • Intra-abdominal Infections • Bacterial Meningitis